
FAUNISTIC RECORDS FROM THE CZECH REPUBLIC – 109

Hymenoptera: Aculeata**Apoidea:****Andrenidae**

Andrena intermedia Thomson, 1870. Bohemia mer., Jankov near České Budějovice (7051), 500 m a.s.l., 17.vii.1933, 1 ♂, L. Baťa leg., J. Straka det., coll. National Museum (Praha); Bohemia or., Hradec Králové - Na Plachtě (5861), 240 m a.s.l., 1.vii.1998, 1 ♂, at locality with sandy soil, J. Straka leg., det. et coll. *Andrena intermedia* was supposed to be a boreoalpine species (Schmid-Egger & Scheuchl 1997). Distribution: Poland, Germany, the Alps (France, Italy) up to 1600 m a.s.l., western and central France, Belgium, Romania, the Czech Republic, Yugoslavia, Bulgaria, Turkey, eastern Sweden and Finland up to 66°N (Dylewska 1987). In the Czech Republic reliably known only from Moravia (Kocourek 1989). Baťa (1952) recorded it from southern Bohemia near České Budějovice; Kocourek (1966) regarded this record as incorrect. Confirmed occurrence in Bohemia.

Colletidae

Hylaeus lepidulus Cockerell, 1924. Bohemia centr.: Praha - Újezd nad Lesy, Blatovská louka (5953), 31.vii.1996, 2 ♀♀, wet meadow; Praha - Hloubětín, Cihelna v Bažantnici (5853), 28.v.1998, 1 ♀; Praha - Horní Počernice, Chvalský lom (5853), 6.vii.1998, 1 ♂, Praha - Hostivař, Toulcův dvůr (5953), 4.vi.1998, 1 ♀; all J. Straka leg. et coll., A. Přidal det. This species was treated as a synonym of *H. gracilicornis* (Morawitz, 1867) (Kocourek 1989) which also lives in the Czech Republic. Distribution: Germany, Austria, Switzerland, Poland, Russia and Mongolia (Dathe et al. 1996). In the Czech Republic known only from Moravia (Přidal 1998). New species for Bohemia.

Hylaeus gracilicornis (Morawitz, 1867). Bohemia centr., Praha – Klánovice, Klánovický les forest (5953), 24.viii.1997, 1 ♀, J. Straka leg. et coll., A. Přidal det. The previous records of this species (e.g., Kocourek 1989) could be based on misidentification (see above under *H. lepidulus* Cockerell, 1924). First reliable record from the Czech Republic.

Pompiloidea: Pompilidae

Agenioideus apicalis (Vander Linden, 1827). Bohemia centr., Praha - Liboc, Divoká Šárka (5951), 27.vi.1998, 1 ♂, rocky steppe, in Moericke's trap, J. Straka leg., det. et coll. Distribution: southern parts of Central Europe and southern Europe, northern Africa (the Atlas Mts.) and Near East (Oehlke & Wolf 1987). In the Czech Republic known only from Moravia (Šedivý 1989, Wolf 1971). New species for Bohemia.

Agenioideus usurarius (Tournier, 1889). Bohemia centr.: Praha - Hloubětín, Cihelna v Bažantnici (5853), 30.ix.1997, 1 ♀, 7.vii.1998, 1 ♂, 1 ♀, 2.viii.1998, 6 ♂♂, 4 ♀♀, 17.viii.1998, 3 ♂♂, 4 ♀♀; Praha - Liboc, Divoká Šárka (5951), 18.ix.1998, 1 ♀; Praha - Horní Počernice, Chvalské skály (5853), 6.vii.1998, 1 ♀; Praha - Horní Počernice, Chvalský lom (5853), 28.vi.1998, 1 ♀, 3.viii.1998, 2 ♂♂; at various localities with dry and warm microclimate, in Moericke's traps; all J. Straka leg. et coll., the specimen captured on 30.ix.1997 at Praha - Hloubětín J. v.d. Smissen det., the others J. Straka det. Distribution: from southern parts of Central Europe to Near East (Oehlke & Wolf 1987). There are two old records from the Czech Republic - one from Bohemia and one from Moravia (Wolf 1971). Confirmed occurrence in the Czech Republic.

Evagetus alamannicus (Blüthgen, 1944). Bohemia bor., Jizerské hory Mts., Vápenný vrch near Raspenava (5056), 21.vi.1997, 1 ♀, lime quarry; Bohemia centr., Praha - Troja, on the edge of ZOO garden (5852), 10.vii.1997, 1 ♂; Bohemia or., Velká Stráž (6260), 20.vii.1997, 1 ♀; all J. Straka leg. et coll., J. v.d. Smissen det. Distribution: from southern Central Europe to the Crimea and Caucasus (Oehlke & Wolf 1987). Only one old specimen known from Bohemia (Wolf 1971). Confirmed occurrence in the Czech Republic.

Sphecoidea: Pemphredonidae

Diodontus brevilabris Beaumont, 1967. Bohemia centr.: Praha - Hloubětín, Cihelna v Bažantnici (5853), 7.vii.1998, 1 ♀, 2.viii.1998, 1 ♂, 1.ix.1998, 1 ♀; Praha - Horní Počernice (5853), 29.vii.1997, 1 ♀, 1 ♂, 3.vii.1997, 1 ♀, 4.vii.1997, 2 ♂♂, 17.vii.1997, 1 ♀; most specimens in Moericke's traps; all J. Straka leg., det. et coll. Distribution: Armenia, Azerbaijan, Bulgaria, Cyprus, Greece, Hungary, Israel, Lebanon, Russia, Turkey, Turkmenistan and Ukraine (Budrys 1998). New species for the Czech Republic.

Spilomena mocsaryi Kohl, 1898. Bohemia centr.: Praha - Troja, Sklenářka (5852), 28.vi.1998, 1 ♀, grassy steppe; Praha - Horní Počernice, Chvalské skály (5853), 18.viii.1998 and 2.ix.1998, 2 ♀♀ captured and 2 ♀♀ found in Moericke's trap, on loess wall; all J. Straka leg., det. et coll. Distribution: Central and southern Europe (Dollfuss 1991). In the Czech Republic known only from Moravia (Pádr 1989a). New species for Bohemia.

Vespoidea:

Eumenidae

Ancistrocerus auctus (Fabricius, 1793). Bohemia centr.: Praha - Zlíchov (5952), 1.vii.1908, 1 ♀, O. Šustera leg., J. Straka det., coll. National Museum (Praha); Praha - Horní Počernice, Chvalský lom (5853), 28.vi.1997, 1 ♀, at locality with sandy soil and loess walls, J. Straka leg., det. et coll. Distribution: from south-eastern Europe to Moravia and Slovakia in the north and Central Asia in the east (Gusenleitner 1995). In the Czech Republic known only from Moravia (Pádr 1989b). New species for Bohemia.

Eumenes sareptanus insolatus Müller, 1923. Bohemia centr.: Kutná Hora (6057), 17.viii.1921, 1 ♀, V. Zavadil leg., J. Gusenleitner det., J. Straka revid.; Kolín (5957), 22.vii.1925, 1 ♂, V. Zavadil leg., J. Gusenleitner det.; Praha - Hlubočepy (5952), ix.1954, 1 ♀, Kočmíd leg., J. Straka det.; Káraný nad Labem (5854), without date, 4 ♂♂, 4 ♀♀, Obenberger leg., J.

Gusenleitner det., J. Straka revid.; all coll. National Museum (Praha). In the Czech Republic known only from Moravia (Pádr 1989b). New species for Bohemia.

Vespidae

Polistes bischoffi Weyrauch, 1937. Moravia mer., Bílé Karpaty Mts., Vápenky settlement (7171b), 2.ix.1998, 1 ♂, J. Strejček leg., J. Straka det. et coll., 23.viii.1999, 15 ♂♂, J. Straka leg., det. et coll. Distribution: southern Germany, Austria, Switzerland, Spain, France, Greece, Yugoslavia and the former USSR (Schmid-Egger 1989). New species for the Czech Republic.

BATĀ L. 1952: Nové nálezy vzácnějších blanokřídlých v jižních Čechách. (De novis rariorum Hymenopterorum speciebus in Bohemia meridionali inventis). *Časopis Čs. Společ. Entomol.*, 49: 165-169 (in Czech, Russian and Latin abstr.). – BUDRYS E. 1998: Two 'rare' wasps, *Diodontus brevilabris* and *Polemistus abnormis* (Hymenoptera, Sphecidae), from south-west Russia. *Acta Zoologica Lituanica*, 8(3): 81-85. – DATHE H. H., HEIDE A. & WITT R. 1996: Nachweis einer neuen Maskenbiene für Europa – *Hylaeus lepidulus* Cockerell, 1924. *Entomol. Nachr. Ber.* (Dresden), 40: 157-163. – DYLEWSKA M. 1987: Die Gattung *Andrena* F. in Nord- und Mitteleuropa. *Acta Zool. Cracov.*, 30: 359-708. – DOLLFUSS H. 1991: Bestimmungsschlüssel der Grabwespen Nord- und Zentraleuropas (Hymenoptera, Sphecidae) mit speziellen Angaben zur Grabwespenfauna Österreichs. *Stapfia*, 24: 1-247. – GUSENLEITNER J. 1995: Bestimmungstabellen mittel- und südeuropäischer Eumeniden (Vespoidea, Hymenoptera). Teil 4: Die Gattung *Ancistrocerus* Wesmäl 1836 mit einem Nachtrag zum Teil 1: Die Gattung *Leptochilus* Saussure. *Linzer Biol. Beitr.*, 27: 753-775. – KOCOUREK M. 1966: Prodrómus insectorum Bohemoslovakiae. Apoidea, *Andrena*. Prodrómus der Hymenopteren der Tschechoslowakei. Pars 9: Apoidea, 1. *Acta Faun. Entomol. Mus. Nat. Pragae*, 12 (Suppl. 2): 3-122. – KOCOUREK M. 1989: Apoidea, pp. 173-184. In: ŠEDIVÝ J. (ed.) 1989: Enumeratio insectorum bohemoslovakiae. Check-list of Czechoslovak insects III (Hymenoptera). *Acta Faun. Entomol. Mus. Nat. Pragae*, 19: 1-195. – OEHLKE J. & WOLF H. 1987: Beiträge zur Insekten-Fauna der DDR: Hymenoptera – Pompilidae. *Beitr. Entomol.* (Berlin), 37: 279-390. – PÁDR Z. 1989a: Sphecoidea, pp. 165-171. In: ŠEDIVÝ J. (ed.) 1989: Enumeratio insectorum bohemoslovakiae. Check-list of Czechoslovak insects III (Hymenoptera). *Acta Faun. Entomol. Mus. Nat. Pragae*, 19: 1-195. – PÁDR Z. 1989b: Vespoidea, pp. 157-160. In: ŠEDIVÝ J. (ed.) 1989: Enumeratio insectorum bohemoslovakiae. Check-list of Czechoslovak insects III (Hymenoptera). *Acta Faun. Entomol. Mus. Nat. Pragae*, 19: 1-195. – PÁDR Z. 1990: Studie výskytu žahadlových blanokřídlých (Hymenoptera – Aculeata) na území Prahy. (Das Vorkommen der Stachelhautflügler (Hymenoptera – Aculeata) in Prag, der Hauptstadt der Tschechoslowakei). *Natura Pragensis*, 7: 1-179 (in Czech, German summary). – PRIDAL A. 1998: New records and additional notes on faunistics of solitary bees (Hymenoptera: Apoidea) from Czech and Slovak Republic. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis*, 46: 27-31. – SCHMID-EGGER Ch. & SCHEUCHL E. 1997: *Illustrierte Bestimmungstabellen der Wildbienen Deutschlands und Österreichs unter Berücksichtigung der Arten in der Schweiz. Band 3: Andrenidae*. Eigenverlag, Velden/Vils, 180 pp. – SCHMID-EGGER Ch. & TREIBER R. 1989: Die Verbreitung von *Polistes bischoffi* (Weirauch 1937) (Hym., Vespoidea) in Süddeutschland. *Linzer Biol. Beitr.*, 21: 601-609. – ŠEDIVÝ J. 1989: Pompiloidea, pp. 161-164. In: ŠEDIVÝ J. (ed.) 1989: Enumeratio insectorum bohemoslovakiae. Check-list of Czechoslovak insects III (Hymenoptera). *Acta Faun. Entomol. Mus. Nat. Pragae*, 19: 1-195. – WOLF H. 1971: Prodrómus insectorum Bohemoslovakiae. Hymenoptera, Pompiloidea. Prodrómus der Hymenopteren der Tschechoslowakei. Pars 10: Pompiloidea. *Acta Faun. Entomol. Mus. Nat. Pragae*, 14(Suppl. 3): 3-76.

Jakub STRAKA, Náchodská 1680,
CZ-193 00 Praha 9 – Horní Počernice, Czech Republic;
e-mail: straka-jakub@mbox.vol.cz