– SMETANA A. 1964: Systematische und faunistische Beiträgentnis der Staphyliniden-Fauna der Tschechoslowakei VI. (Col., Staphylinidae). *Čas. Čs. Společ. Entomol.*, 61: 162-172. – SMETANA A. 1966: Systematische und faunistische Beiträge zur Kenntnis der Staphylinidenfauna der Tschechoslowakei VII. (Col., Staphylinidae). *Acta Entomol. Bohemoslov.*, 63: 322-336. – SMETANA A. 1993: Staphylininae, pp. 47-50. In JELÍNEK J.: (ed.) Check-list of Czechoslovak Insects IV (Coleoptera). Seznam československých brouků. *Folia Heyrovskyana*, Suppl. 1: 3-172 (in English and Czech).

Jan MATĚJÍČEK, Formánkova 436, CZ-500 11 Hradec Králové 11, Czech Republic
Jaroslav BOHÁČ, Institute of Landscape Ecology, Academy of Sciences of the Czech Republic, Na Sádkách 7, CZ-370 05 České Budějovice, Czech Republic

Published November 26, 2003

Klapalekiana, **39**: 135, 2003
ISSN 1210-6100

FAUNISTIC RECORDS FROM THE CZECH REPUBLIC – 167

Coleoptera: Carabidae

Epaphius rivularis (Gyllenhal, 1810). Moravia mer. occ., 1 km east of Klatovec, National Nature Reserve Zhejral (6757), 680 m a.s.l., 4.vi.-6.viii.2002, 1 ♂, in pitfall trap, J. Kaláb lgt., det. et coll. The trap was set in a peat-bog with solitary single trees and bushes (*Alnus* sp., *Betula* sp., *Juniperus* sp., *Picea* sp., *Pinus* sp., *Salix* sp.) on the unshaded northern shore of the Zhejral pond. Euro-Siberian species; from the Czech Republic so far known only from Bohemia (Bašta et al. 2002), where it is rare and locally distributed in wet habitats from lowland to mountain zones. New species for Moravia.

BAŠTA J., KALÁB J. & MORAVEC P. 2002: Distribution and ecology of Epaphius rivularis (Coleoptera: Carabidae) in the Czech Republic. *Klapalekiana*, 38: 163-171.

Jaroslav KALÁB, Jinačovice 119, CZ-664 34 Kuřim, Czech Republic