Published December 31, 2005

Klapalekiana, **41**: 263-264, 2005
ISSN 1210-6100

FAUNISTIC RECORDS FROM THE CZECH REPUBLIC – 195

Coleoptera:

Curculionidae

Bagous lutulosus (Gyllenhal, 1827). Bohemia or., Hradec Králové, Na Plachtě Nature Monument (5861a), 6.vi.2002, 1 ♂, T. Kopecký leg., J. Krátký det. et coll. Western Palaearctic species living on *Juncus* spp. (Caldara & O'Brien 1997). In former Czechoslovakia reliably known only from Slovakia, its occurrence in Moravia was regarded as uncertain (Strejček 1993). However, the species was already reported by Roubal (1970) from Bohemia (Jesenice, Lázský rybník pond, without date and collector's name). This is the second record from Bohemia and confirmation of the occurrence of this species in the Czech Republic after more than 30 years.

Baris (Baris) atramentaria (Boheman, 1836). Bohemia bor, Raná (hill) (5548d), 24.iii.2002, 1 ♀, R. Škoda leg., det. et coll., J. Krátký revid. Collected by sifting a substrate near plants of *Erysimum* sp. In the Czech Republic known only from Moravia (Strejček 1993). The first record from Bohemia.

Ceutorhynchus subpilosus Ch. Brisout, 1869. Moravia mer., Brno – Hády (6766c), without date, 1 ♀, Stejskal leg., J. Krátký det., coll. Muzeum východních Čech (Hradec Králové). Mediterranean and Caucasian species known also from Slovakia (Strejček 1993). It lives on *Alyssum alyssoides*. New species for the Czech Republic.

Ceutorhynchus plumbeus Ch. Brisout, 1869. Moravia, "Babice" [one of several settlements with this name], iv.1944, 1 ♂, without name of collector, J. Krátký det, coll. Muzeum východních Čech (Hradec Králové); Moravia mer., Brno – Řečkovice (6765c), without date, 1 ♀, Štaif leg., S. Benedikt det., coll. Západočeské muzeum (Plzeň). It lives on *Erysimum cheiranthoides* in sandy habitats. In the Czech Republic known from Bohemia (Strejček 1993). First record from Moravia.

Ceutorhynchus merkli Korotyaev, 2000. Moravia mer.: Čejč – Špidlák (7067d), 7.v.1994, 1 ♂, 23.iv.1995, 3 ♂♂, 9 ♀♀, 12.v.1996, 2 ♀♀, 26.iv.1996, 2 ♀♀, J. Krátký leg., det. et coll.; Čejkovice (7067d), 6.vi.1996, 1 ♀, Z. Ivánek leg., S. Benedikt det. et coll.; Kobylí (7067c), 25.iv.2000, 3 ♂♂, 3 ♀♀, J. Krátký & T. Kopecký leg. et coll., J. Krátký det., 8.v.2003, 2 spec., T. Kopecký leg. et coll., J. Krátký det. South-eastern European species living on *Cardaria draba* (Korotyaev 2000). In the Czech Republic only in southeastern part of Moravia, in many habitats not rare in spring. Previously confused with the closely related *C. ignitus* Germar, 1824, and sometimes also identified as *C. suturellus* Gyllenhal, 1837 (Korotyaev 2000). New species for the Czech Republic.

Datonychus urticae (Boheman, 1845). Moravia or., Vizovice (6773c), 25.v.1944, 1 spec., V. Balthasar leg., J. Fremuth det., coll. Muzeum východních Čech (Hradec Králové). In central Europe a rare species living in forests as well as in steppes on *Stachys* spp. New species for the Czech Republic.

Brachysomus dispar Penecke, 1910. Moravia mer., Moravský kras Karst, Habrůvecká bučina Nature Reserve (6666c), 20.v.2002, 2 ♂♂, 1 ♀, T. Kopecký leg. et coll., S. Benedikt det., 8.vi.2002, 1 ♂, 3 ♀♀, J. Krátký leg., det. et coll. Collected by sifting in an oak and beach forest. Preferring limestone substrate and so far known from the western Carpathians (Slovakia, Ukraine, Poland, and Hungary) (Benedikt 2001). New species for the Czech Republic.

Larinus* (*Larinomesius*) *obtusus Gyllenhal, 1836. Bohemia centr., Semice env., Semická hůra hill (5855a), 12.v.2000, 1 spec., B. Zbuzek leg. et coll., J. Krátký det.; Moravia mer.: Bzenec – Přívoz (7069), 11.vii.1998, 1 spec., L. Koloničný leg. et coll., S. Benedikt det.; Nesyt pond near Mikulov (7266a), 29.v.1999, L. Blažej leg. et coll., S. Benedikt det. Expansive species living on *Centaurea* spp. In former Czechoslovakia known only from Slovakia (Strejček 1993). New species for the Czech Republic.

Nanophyidae

Dieckmanniellus helveticus (Tournier, 1867). Moravia mer., Břeclav – Pohansko (7267d), 25.vii.2000, 1 ♂, 1 ♀, J. Krátký leg. et coll., J. Fremuth det. Collected by sweeping near a water channel. In the Czech Republic known only from Bohemia (Strejček 1993). Its occurrence in Moravia has been expected with regard to its occurrence in Bohemia, and is thereby confirmed.

BENEDIKT S. 2001: Brachysomus rokosensis sp. n. from Slovakia with remarks on distribution and bionomics of the genus Brachysomus in the Western Carpathian Mts. (Coleoptera: Curculionidae). *Klapalekiana*, 37: 139-146. – CALDARA R. & O'BRIEN Ch. W. 1997: Systematics and evolution of weevils of the genus Bagous VI. Taxonomic treatment of the species of the western Palearctic Region (Coleoptera: Curculionidae). *Mem. Soc. Entomol. Ital.*, 76: 131-347. – KOROTYAEV B. A. 2000: A new species of Ceutorhynchus Germar, 1824 living on Cardaria draba in southeastern Europe (Coleoptera, Curculionidae). *Acta Zoologica Academiae Scientiarum Hungaricae*, 46(4): 305-308. – SMRECZYŃSKI S. 1974: *Podrodzina Ceutorhynchinae. Klucze do oznaczania owadów Polski. Cz. XIX: Coleoptera, Zesz. 98e: Curculionidae.* PWN, Warszawa, 180 pp (in Polish). – STREJČEK J. 1993: Curculionidae, pp. 135-152. In: JELÍNEK J. (ed.): Check list of Czechoslovak Insects IV (Coleoptera). Seznam československých brouků. *Fol. Heyrovskyana*, Suppl. 1: 3-172 (in English and Czech). – ROUBAL J. 1970: Entomologický průzkum Příbramska. *Vlastivědný Sborník Podbrdska*, 1970: 156-187 (in Czech).

Jiří KRÁTKÝ, Třebechovická 821, CZ-500 03 Hradec Králové, Czech Republic