

FAUNISTIC RECORDS FROM THE CZECH REPUBLIC – 230

Coleoptera: Trogidae

Trox perrisii Fairmaire, 1868. Bohemia centr., Lány – “Kouglouva ohrádka” (5849D21) [= locality No. 8: “Lánská obora – Kouglův rybník”, fig. 1, for details see Januš 2004: 60], 18.vi.2002, 1 spec., at light, K. Rébl lgt., det. et coll., D. Král revid.; the only specimen collected from forest pasture covered with several old solitary trees of oak (*Quercus*) together with two specimens of *Trox scaber* (Linnaeus, 1767). A nidicolous species collected mostly from nests in tree hollows e. g., of hoopoes (*Upupa epops*) or from “free” nests of herons (*Ardea*) or storklings (*Ciconia*) (Král & Vitner 1993, Martín-Piera & López-Colón 2000). Distributed in Northern Africa (Algeria, Morocco), western parts of Europe and in Turkey; in Central Europe rarely collected in Austria, Czech Republic, Hungary, Germany, Poland and Slovakia (cf. e. g., Baraud 1985, 1992; Martín-Piera & López-Colón 2000; Pittino 2006); recent records are e. g., from Germany from Berlin (Esser 2002) and from the Rhine province (Hadulla 2003). In the Czech Republic previously known from southern Moravia only, these findings and findings from Slovakia summarized by Král & Vitner (1993) and Juřena et al. (2000). New species for Bohemia.

BARAUD J. 1985: *Coléoptères Scarabaeoidea. Faune du Nord de l'Afrique du Maroc au Sinai. Encycl. Entomol.*, 46. Editions Lechevalier, Paris, 651 pp. – BARAUD J. 1992: *Coléoptères Scarabaeoidea d'Europe. Faune de France* 78. Féd. Fr. Soc. Entomol. & Soc. Linn. de Lyon, Lyon, 856 pp. – ESSER J. 2002: Neue Käferfunde aus Berlin (Teil 3). *Märkische Entomol. Nachr.*, 4: 45–47. – HADULLA K. 2003: Bemerkungen zu einigen Neu- und Wiederfunden für die Rheinprovinz (Col., Trogidae, Scarab., Curculion.). *Mitt. Arbeitsgem. Rheinischer Koleopterol.*, 13: 40–42. – JANUŠ J. 2004: Výsledky faunistického inventarizačního průzkumu brouků (Coleoptera) čeledi Chrysomelidae s. lat. na území Chráněné krajinné oblasti a Biosférické rezervace Křivoklátsko. (Results of a survey of the beetles (Coleoptera) of family Chrysomelidae s. lat. in the territory of the Křivoklátsko Protected Landscape Area and Biosphere Reserve). *Klapalekiana*, 40: 55–121 (in Czech, English summary). – JUŘENA D., BEZDĚK A. & TÝR V. 2000: Zajímavé nálezy listorohých brouků (Coleoptera: Scarabaeoidea) na území Čech, Moravy a Slovenska. (Interesting faunistic records of Scarabaeoidea (Coleoptera) from Bohemia, Moravia and Slovakia). *Klapalekiana*, 36: 233–257 (in Czech, English summary). – KRÁL D. & VITNER J. 1993: Faunistické síťové mapování listorohých brouků (Coleoptera: Scarabaeoidea) Československa – výběr výsledků získaných v letech 1989–1990. (Faunistic grid mapping of Czechoslovak Scarabaeoidea (Coleoptera) – selected results obtained in 1989–1990). *Klapalekiana*, 29: 25–36 (in Czech, English summary). – MARTÍN-PIERA F. & LÓPEZ-COLÓN J. I. 2000: *Coleoptera, Scarabaeoidea I: Fauna Ibérica, vol. 14*. Museo Nacional de Ciencias Naturales, Consejo Superior de Investigaciones Científicas, Madrid, 526 pp. – PITTINO R. 2006: Family Trogidae MacLeay, 1819, pp. 79–81. In: LÖBL I. & SMETANA A. 2006: *Catalogue of Palaearctic Coleoptera, Vol. 3. Scarabaeoidea – Scirtoidea – Dasciloidea – Buprestoidea – Byrrhoidea*. Apollo Books, Stenstrup, 690 pp.

David KRÁL, Department of Zoology, Faculty of Science, Charles University in Prague,
CZ-128 44 Praha 2, Czech Republic; e-mail: kraldavid@centrum.cz
Karel RÉBL, Žižkovo náměstí 976, CZ-271 01 Nové Strašecí, Czech Republic;
e-mail: karel.rebl@zukolin.cz