
FAUNISTIC RECORDS FROM THE CZECH REPUBLIC – 324

Coleoptera: Staphylinidae:**Micropeplinae**

Micropeplus latus C. Hampe, 1861. Bohemia mer.: Šumava mts., Volary env., Boubín National Nature Reserve (7048), 25.xi.2006, 1 ♂, in flight in primary forest at a temperature of ca. 5°C, J. Vávra leg. and coll., J. Vávra and P. Štourač det. Little known and extremely rare, probably a relict species. Distributed discontinuously in Europe, reported only from Croatia, Greece, Slovakia and Sweden (Smetana 2004b), and also from Slovenia (“Krain”) by Horion (1963), Austria (Niederösterreich) (Kapp 2003), France and Italy (Tronquet 2008). New species for the Czech Republic.

Proteininae

Megarhtrus stercorarius Mulsant et Rey, 1878. Moravia bor.: Hrubý Jeseník mts., Velká kotlina valley (5969), 1200 m a.s.l., 26.viii.2006, 2 ♂♂, sifted from rotten grass in meadow near spring, M. Mantič (Hlučín-Bobrovniky, Czech Republic) leg., det. and coll.; Hrubý Jeseník mts., Ovčárna challet env., Velký Dědův důl valley (5969), 1270 m a.s.l., 6.viii.2010, 2 ♂♂, in old fruiting bodies of *Russula* sp., J. Vávra leg., det. and coll.; same locality, 1300 m a.s.l., 11.ix.2010, 3 ♂♂, 2 ♀♀, in excrement of Cervidae, J. Vávra leg., det. and coll. Moravia bor. or.: Moravskoslezské Beskydy mts., Staré Hamry-Břestovy (6576), 600 m a.s.l., 2.v.1998, 1 ♂, sifted near manger with old hay in Picetum, M. Mantič leg., det. and coll. European species known only from Austria, Bulgaria, Croatia, Georgia, Italy, Russia: South European Territory and Switzerland (Smetana 2004a). New species for the Czech Republic.

Oxytelinae

Bledius (Hesperophilus) secessus Bondroit, 1912. Moravia mer.: Mikulov env., Svatý kopeček hill (6175), 6.vii.1984, 1 spec., gravel in a calcareous quarry, P. Štourač (Praha, Czech Republic) leg., det. and coll.; same locality, 19.iv.2007, 7 spec., in earth wall of a quarry, J. Vávra leg., M. Schülke det., coll. J. Vávra and M. Schülke (Berlin, Germany); same locality, 11.v.2008, 2 spec., T. Sitek (Ostrava, Czech Republic) leg. and coll., J. Vávra det. Species known only from Croatia, Germany and Italy (Smetana 2004e). New species for the Czech Republic.

Steninae

Stenus cribratus Kiesenwetter, 1850. Moravia mer.: Zaječí env., Kalvárie hill (7166), 17.x.2002, 1 ♂, under stone on steppe, J. Vávra leg., det. and coll. Widespread throughout southern Europe, known also from southern part of central Europe and Asia (Turkey). In central Europe reported only from Austria, Hungary and Slovakia (Smetana 2004f). New species for the Czech Republic.

Staphylininae

Bisnius nigriventris (C. G. Thomson, 1867). Moravia bor. or.: Moravskoslezské Beskydy mts., Dolní Lomná env., Mionší National Nature Reserve (6478), 850 m a.s.l., 16.vii.2006, 1 ♂, in old *Polyporus squamosus* on trunk of *Fagus sylvatica*, J. Vávra leg., det. and coll. Euro-Siberian species, in central Europe known from Austria, Czech Republic, Hungary and Switzerland (Smetana 2004g), and also from Germany (Köhler & Klausnitzer 1998), Poland (Szujecki 1995) and Slovakia (Dvořák 1967). In the Czech Republic known only from Lokay's old data from Bohemia reported by Smetana (1958) and from several localities in eastern Bohemia reported by Jelínek & Voda (1999). New species for Moravia.

Tachyporinae

Bryoporus multipunctus Hampe, 1867. Moravia mer.: Vranov nad Dyjí-Braitava forest (7160–7161), 24.v.2007, 2 spec., swept in deciduous forest, J. Vávra leg. and coll.; Vranov nad Dyjí-Ledové sluje (7161), 23.v.2007, 3 spec., swept in mixed forest, J. Vávra leg., coll. J. Vávra and P. Štourač. All specimens P. Štourač det. Widespread throughout southern and south-eastern Europe, known also from Asia (Israel, Lebanon and Turkey) (Schülke 2003, Smetana 2004c). In central Europe known from Austria, Hungary, Slovakia and Switzerland (Schülke 2003), and also from Poland (Pašník 1998). From the Czech Republic reported only in Catalogue of Palaearctic Coleoptera by Smetana (2004c), but without precise data. First concrete record from the Czech Republic.

Aleocharinae

Atheta (Atheta) ebenina (Mulsant et Rey, 1873). Moravia bor.: Hrubý Jeseník mts., Velká kotlina valley (5969), 2.viii.2007, 1 ♀, J. Vávra leg. and coll., P. Štourač det. European species, in central Europe known from Austria, Czech Republic, Germany, Poland, Slovakia and Switzerland (Smetana 2004d). In the Czech Republic reported from Bohemia ("Sudety") by Fleischer (1930); from Moravia a dubious record by Likovský (1993). First certain record from Moravia.

Atheta (Microdota) excelsa Bernhauer, 1911. Moravia bor.: Hrubý Jeseník mts., Ovčárna challet (5969), 1150 m a.s.l., 12.ix.2008, 1 ♂, sifted in Picetum and near brook, J. Vávra leg. and coll., P. Štourač det. Euro-Siberian species, in central Europe known from Austria, Germany, Poland, Slovakia and Switzerland (Smetana 2004d). New species for the Czech Republic.

Atheta (Oreostiba) subglabra (Sharp, 1869). Moravia bor.: Hrubý Jeseník mts., Velká kotlina valley (5969), 1100–1200 m a.s.l., 13.vi.2007, 1 ♀, J. Vávra leg. and coll., P. Štourač det. Species distributed in western, northern and central Europe. In central Europe known from Austria, Germany, Poland and Slovakia (Smetana 2004d). New species for the Czech Republic.

Ischnoglossa elegantula (Mannerheim, 1830). Moravia bor.: Hrubý Jeseník mts., Ovčárna challet (5969), 27.viii.2000, 1 ♂, 1 ♀, J. Vávra leg., S. Snäll det., P. Wunderle revid., coll. J. Vávra and P. Wunderle (Mönchengladbach, Germany); Hrubý Jeseník mts., Ovčárna challet

env., Velký Dědův důl valley (5969), 1200–1280 m a.s.l., 6.viii.2010, 3 ♂♂, 2 ♀♀; same locality, 9.viii.2010, 3 ♂♂, 3 ♀♀; same locality, 11.ix.2010, 1 ♂; all J. Vávra leg., det. and coll. All specimens collected under bark of old trunks of *Picea abies*. European species known from Finland, Italy, Lithuania, Norway, Russia: North European Territory and Sweden (Smetana 2004d), and also from Albania, Austria, Slovenia and Switzerland (Wunderle 1990). New species for the Czech Republic.

Ischnoglossa obscura Wunderle, 1990. Moravia bor.: Hrubý Jeseník mts., Vidly env., Jelení bučina Nature Reserve (5869), 28.xi.2009, 1 ♂, sifted in mixed forest, J. Vávra leg., det. and coll. Moravia bor. or.: Moravskoslezské Beskydy mts., Ostravice (6476), 25.viii.1990, 1 ♂, 1 ♀, J. Vávra leg., P. Štourač det., P. Wunderle revid., coll. J. Vávra and P. Wunderle. Species known from Europe and also from Asia (Turkey) (Smetana 2004d, Assing 2007). In Europe reported only from Austria, Belgium, France, Germany and Sweden (Smetana 2004d), and also from Denmark (Hansen 1996). New species for the Czech Republic.

Aleochara (Xenochara) helvetica Likovský, 1982. Moravia bor.: Hrubý Jeseník mts., Velká kotlina valley (5969), 1150 m a.s.l., 3.vi.2008, 10 spec., J. Vávra leg. and det., coll. J. Vávra, S. Snäll (Tumba, Sweden) and P. Štourač; same locality, 1200 m a.s.l., 26.v.2009, 4 spec., J. Vávra leg., det. and coll. All specimens were collected in the alpine zone close to snow, together with the related species *Aleochara heeri* Likovský, 1982, and with *Deliphrosoma prolongatum* (Rottenberg, 1873), *Quedius alpestris* (Heer, 1839) and *Q. unicolor* Kiesenwetter, 1848 (all Staphylinidae). Species known only from the Alps in Austria, Germany and Switzerland (Smetana 2004d). New species for the Czech Republic.

ACKNOWLEDGEMENTS. I would like to thank M. Mantič (Hlučín-Bobrovníky, Czech Republic), T. Sitek (Ostrava, Czech Republic) and P. Štourač (Praha, Czech Republic) for providing additional data for *Megarthritis stercorarius* and *Bledius secessus*; M. Schülke (Berlin, Germany), P. Štourač and P. Wunderle (Mönchengladbach, Germany) for identification of some species. V. Assing (Hannover, Germany) and A. Byk (Warszawa, Poland) are thanked for help with obtaining literature, and J. Janák (Rtyně nad Bílinou, Czech Republic) and J. Růžicka (Praha, Czech Republic) for critical comments on the manuscript. M. V. L. Barclay (London, Great Britain) is acknowledged for revision of English.

ASSING V. 2007: New species and additional records of Staphylinidae from Turkey V (Coleoptera). *Stuttgarter Beiträge zur Naturkunde, Serie A (Biologie)* 700: 1–64. – DVOŘÁK M. 1967: Zajímavé nálezy drabčůků na Slovensku II (Col., Staphylinidae). (Interessante Funde der Staphyliniden-Arten aus der Slowakei II (Col., Staphylinidae). *Acta Rerum Naturalium Musei Nationalis Slovenici, Bratislava* 13(1): 81–88 (in Czech, German summary). – FLEISCHER A. 1930: Přehled brouků fauny Československé republiky. [Overview of the beetle fauna of the Czechoslovak Republic]. Moravské museum zemské, Brno, 485 pp. (in Czech). – HANSEN M. 1996: Katalog over Danmarks biller. Catalogue of the Coleoptera of Denmark. *Entomologiske Meddelelser* 64(1–2): 1–231 (in Danish and English). – HORION A. 1963: Faunistik der mitteleuropäischen Käfer. Bd. 9. Staphylinidae. 1. Teil. Micropeplinae bis Euaesthetinae. A. Feyel, Überlingen-Bodensee, xi+412 pp. – JELÍNEK J. & VODA J. 1999: Drabčici Orlických hor a Podorlicka (Coleoptera, čeled' Staphylinidae), podčeledi Staphylininae, tribu Philonthini. [Rove beetles of the tribe Philonthini of the subfamily Staphylininae from Orlické hory mountains and Podorlicko region (Coleoptera, Staphylinidae)]. *Orlické hory a Podorlicko* 9: 31–44 (in Czech). – KAPP A. 2003: 2216. Micropeplus latus Hampe, 1861 – in Österreich (Staphylinidae). *Entomologische Blätter* 98: 178. – KÖHLER F. & KLAUSNITZER B. (eds) 1998: Verzeichnis der Käfer Deutschlands. *Entomologische Nachrichten und Berichte* (Dresden), Beiheft 4: 1–185. – LIKOVSKÝ Z. 1993: Staphylinidae: Aleocharinae. Pp. 52–62. In: JELÍNEK J. (ed.): Check-list of Czechoslovak Insects IV (Coleoptera). Seznam československých brouků. *Folia Heyrovskyana, Supplementum* 1: 3–172 (in English and Czech). – PAŠNÍK G. 1998: Bryoporus multipunctus Hampe, 1867 (Coleoptera: Staphylinidae: Tachyporinae), nowy dla fauny Polski przedstawiciel chrząszczy. (Bryoporus multipunctus Hampe, 1867 (Coleoptera: Staphylinidae:

Tachyporinae), a beetle new to the fauna of Poland). *Wiadomości Entomologiczne* **16**[1997]: 155–156 (in Polish, English summary). – SCHÜLKE M. 2003: Zur Taxonomie und Verbreitung westpaläarktischer Arten der Gattungen *Bryoporus* Kraatz und *Mycetoporus* Mannerheim (Coleoptera, Staphylinidae, Tachyporinae). 49. Beitrag zur Kenntnis der Tachyporinen. *Entomologische Blätter* **99**: 27–54. – SMETANA A. 1958: *Fauna ČSR. Svazek 12. Drabčíkoviti – Staphylinidae I. Staphylininae*. Nakladatelství Československé Akademie Věd, Praha, 435 pp. (in Czech, German and Russian summary). – SMETANA A. 2004a: Staphylinidae: Proteininae. Pp. 268–270. In: LÖBL I. & SMETANA A. (eds): *Catalogue of Palaearctic Coleoptera, Vol. 2: Hydrophiloidea – Histeroidea – Staphylinidea*. Apollo Books, Stenstrup, 942 pp. – SMETANA A. 2004b: Staphylinidae: Micropeplinae. Pp. 271–272. In: LÖBL I. & SMETANA A. (eds): *Catalogue of Palaearctic Coleoptera, Vol. 2: Hydrophiloidea – Histeroidea – Staphylinidea*. Apollo Books, Stenstrup, 942 pp. – SMETANA A. 2004c: Staphylinidae: Tachyporinae. Pp. 330–352. In: LÖBL I. & SMETANA A. (eds): *Catalogue of Palaearctic Coleoptera, Vol. 2: Hydrophiloidea – Histeroidea – Staphylinidea*. Apollo Books, Stenstrup, 942 pp. – SMETANA A. 2004d: Staphylinidae: Aleocharinae. Pp. 353–494. In: LÖBL I. & SMETANA A. (eds): *Catalogue of Palaearctic Coleoptera, Vol. 2: Hydrophiloidea – Histeroidea – Staphylinidea*. Apollo Books, Stenstrup, 942 pp. – SMETANA A. 2004e: Staphylinidae: Oxytelinae. Pp. 511–535. In: LÖBL I. & SMETANA A. (eds): *Catalogue of Palaearctic Coleoptera, Vol. 2: Hydrophiloidea – Histeroidea – Staphylinidea*. Apollo Books, Stenstrup, 942 pp. – SMETANA A. 2004f: Staphylinidae: Steninae. Pp. 537–564. In: LÖBL I. & SMETANA A. (eds): *Catalogue of Palaearctic Coleoptera, Vol. 2: Hydrophiloidea – Histeroidea – Staphylinidea*. Apollo Books, Stenstrup, 942 pp. – SMETANA A. 2004g: Staphylinidae: Staphylininae. Pp. 624–698. In: LÖBL I. & SMETANA A. (eds): *Catalogue of Palaearctic Coleoptera, Vol. 2: Hydrophiloidea – Histeroidea – Staphylinidea*. Apollo Books, Stenstrup, 942 pp. – SZUJECKI A. 1995: Zgrupowania kusakowatych (Col., Staphylinidae s. l.) borów sosnowych świeżych i ich antropogeniczne przeobrażenia. [Assemblages of rove beetles (Col., Staphylinidae s. l.) in pine forests and their anthropogenic transformations]. Pp. 175–251. In: SZUJECKI A. (ed.): *Antropogeniczne przeobrażenia epigeicznej i glebowej entomofauny borów sosnowych*. [Anthropogenic transformations of the epigeic and soil entomofauna of pine forests]. Fundacja “Rozwój Szkoły Głównej Gospodarstwa Wiejskiego”, Warszawa, 460 pp. (in Polish). – TRONQUET M. 2008: Staphylinidae Micropeplinae de la faune de France et contrées voisines des genres *Arrhenopeplus* (Koch, 1937) et *Micropeplus* Latreille, 1809 (23^{me} contribution à la connaissance des Staphylins). *Revue de l'Association Roussillonnaise d'Entomologie* **16**[2007]: 104–116.

Jiří Ch. VÁVRA, Ostrava Museum, Masarykovo náměstí 1,
CZ-702 00 Ostrava 1, Czech Republic; e-mail: jiri.vavra@ostmuz.cz