
FAUNISTIC RECORDS FROM THE CZECH REPUBLIC – 347

Diptera: Chironomidae

Paramerina divisa (Walker, 1856). Bohemia mer.: Šebířov (6454), Slupský potok (brook), 49°33'44"N, 14°49'14"E, ca. 405 m a.s.l., 29.v.2012, 1 pupa, N. Lapšanská lgt., I. Skála det. et coll. West Palaearctic species (Sæther & Spies 2013). New species for the Czech Republic.

Cricotopus (Cricotopus) annulator Goetghebuer, 1927. Bohemia sept.: Postoloprty (5648), Ohře (river), 50°20'53"N, 13°41'36"E, ca. 185 m a.s.l., 16.ix.2008, 1 pupal exuvia; Bohemia centr.: Lužec nad Vltavou (5652), Vltava (river), 50°19'09"N, 14°23'53"E, ca. 158 m a.s.l., 12.vii.2012, 3 pupal exuviae; Praha-Podolí (5952), Vltava (river), 50°03'12"N, 14°24'55"E, ca. 187 m a.s.l., 13.vii.2012, 1 pupal exuvia; Praha-Strnady (6052), Vltava (river), 49°56'33"N, 14°23'40"E, ca. 189 m a.s.l., 13.vii.2012, 2 pupal exuviae, all I. Skála lgt., det. et coll. Holarctic species (Sæther & Spies 2013). In the Czech Republic only known from Moravia (Syrůvka et al. 2009). New species for Bohemia.

Cricotopus (Cricotopus) curtus Hirvenoja, 1973. Bohemia centr.: Soutice (6256), Želivka (river), downstream of the Švihov Reservoir, 49°43'50"N, 15°03'18"E, ca. 320 m a.s.l., 27.ix.2011, 1 pupa; Zruč nad Sázavou (6256), Ostrovský potok (brook), 49°44'40"N, 15°06'24"E, ca. 327 m a.s.l., 2.x.2012, 2 pupal exuviae; Hrachov (6351), Brzina (brook), 49°40'12"N, 14°17'35"E, ca. 272 m a.s.l., 21.ix.2011, 1 pupal exuvia; Bohemia or.: Světlá nad Sázavou (6358), Sázavka (brook), 49°40'15"N, 15°24'41"E, ca. 390 m a.s.l., 9.x.2012, 1 pupal exuvia, all I. Skála lgt., det. et coll. Holarctic species (Sæther & Spies 2013). New species for the Czech Republic.

Cricotopus (Cricotopus) triannulatus (Macquart, 1826). Bohemia centr.: Lužec nad Vltavou (5652), Vltava (river), 50°19'09"N, 14°23'53"E, ca. 158 m a.s.l., 12.vii.2012, 1 pupal exuvia; Praha-Troja (5852), Vltava (river), 50°06'51"N, 14°24'52"E, ca. 180 m a.s.l., 13.vii.2010, 4 pupae, 12.vii.2012, 1 pupa; Praha-Dolní Počernice (5953), Rokytka (brook), downstream of the Počernický Pond, 50°05'19"N, 14°34'35"E, ca. 225 m a.s.l., 20.ix.2011, 3 pupae; Píkovice (6152), Sázava (river), 49°52'43"N, 14°25'36"E, 201 m a.s.l., 23.ix.2010, 2 pupal exuviae; Bohemia or.: Dobronín (6559), Mlýnský potok (brook), downstream of the Nový štocký Pond, 49°28'59"N, 15°37'48"E, ca. 470 m a.s.l., 17.v.2011, 1 pupa; Bohemia mer.: Soběslav (6754), Černovický potok (brook), downstream of the Nový u Soběslavi Pond, 49°15'31"N, 14°43'20"E, ca. 403 m a.s.l., 15.x.2012, 6 pupal exuviae, all I. Skála lgt., det. et coll. Holarctic and Neotropical species (Sæther & Spies 2013). In the Czech Republic only known from Moravia (Brabec & Langton 2006). New species for Bohemia.

Cricotopus (Isocladus) intersectus (Staeger, 1839). Bohemia centr.: Praha-Podolí (5952), Vltava (river), 50°03'12"N, 14°24'55"E, ca. 187 m a.s.l., 13.vii.2012, 2 pupal exuviae, 7 pupae, I. Skála lgt., det. et coll. Holarctic species (Sæther & Spies 2013). In the Czech Republic only known from Moravia (Brabec & Langton 2006). New species for Bohemia.

Orthocladius (Symposiocladius) ruffoi Rossaro et Prato, 1991. Bohemia mer.: Tur-ovec (6654), Turovecký potok (brook), downstream of the Turovecký Pond, 49°23'31"N, 14°44'52"E, ca. 410 m a.s.l., 4.v.2010, 1 pupa, I. Skála lgt., det. et coll. Distributed in Europe and North Africa (Sæther & Spies 2013). New species for the Czech Republic.

Chironomus (Lobochironomus) dissidens Walker, 1856. Bohemia centr.: Všebořice (6256), pond northwest of village, 49°42'02"N, 15°08'29"E, ca. 386 m a.s.l., 4.vi.2012, 7 pupal exuviae; Keblov (6356), pond east of village, 49°40'48"N, 15°05'06"E, ca. 394 m a.s.l., 4.vi.2012, 2 pupal exuviae, all I. Skála lgt., det. et coll. Widespread in Europe (Langton & Visser 2003). New species for the Czech Republic.

BRABEC K. & LANGTON P. H. 2006: Faunistic records. Chironomidae. Pp. 143–145. In: KINKOROVÁ J. (ed.): *Dipterologica bohemoslovaca* Vol. 13. *Acta Universitatis Carolinae Biologica* **50**: 1–158. – LANGTON P. H. & VISSER H. 2003: *Chironomidae exuviae. A key to pupal exuviae of the West Palaearctic Region*. CD-ROM, Expert Center for Taxonomic Identification, University of Amsterdam, Amsterdam. – SÆTHER O. A. & SPIES M. 2013: Fauna Europaea: Chironomidae. In: BEUK P. & PAPE T. (eds): *Fauna Europaea: Diptera*. Fauna Europaea version 2.5, <http://www.faunaeur.org> (accessed 11 April 2013). – SYROVÁTKA V., SCHENKOVÁ J. & BRABEC K. 2009: The distribution of chironomid larvae and oligochaetes within a stony-bottomed river stretch: the role of substrate and hydraulic characteristics. *Fundamental and Applied Limnology, Archiv für Hydrobiologie* **174**: 43–62.

Ivan SKÁLA, Povodí Vltavy, s. p. (Vltava River Authority),
Na Hutmance 5a, CZ-158 00 Praha 5-Jinonice, Czech Republic;
e-mail: ivan.skala@pvl.cz, ivan.skala@centrum.cz