Published December 27, 2019 Klapalekiana, **55**: 269–272, 2019
ISSN 1210-6100

FAUNISTIC RECORDS FROM THE CZECH REPUBLIC – 479

Coleoptera: Latridiidae:

Latridiinae

Enicmus amici Lohse, 1981. Moravia mer.: Znojmo, Riegrova ulice street (7162), 10.ix.2017, 1 ♂, beaten from old branches laying in compost, R. Stejskal leg., det. et coll., P. Průdek revid.; Znojmo, Leska Horní street (7162), 30.v.2018, 2 ♀♀, beaten from dry *Verbascum*, R. Stejskal leg. et coll., P. Průdek det.; Znojmo-Oblekovice (7162), 7.vii.2018, 2 ♂♂, 1 ♀, sifted from heap of semidry weeds and grass near garden allotments, R. Stejskal leg. et det., P. Průdek revid., coll. R. Stejskal and Jihomoravské muzeum ve Znojmě, Czech Republic.

West Palaearctic species known from Austria, Bulgaria, Germany, Italy, Poland, Slovakia, Ukraine and Turkey (Rücker 2018). New species for the Czech Republic.

Melanophthalma (***Cortilena***) ***fuscipennis*** (Mannerheim, 1844). Moravia mer.: Podyjí National Park: Lukov – Gálišská louka meadow (7161), 11.ix.2018, 1 ♀, sweeping on dry meadow, R. Stejskal leg., det. et coll.; Podmolí – Šobes (7161), 4.viii.2016, 2 ♀♀, sweeping on forest-steppe, R. Stejskal leg. et coll., P. Průdek det.

West Palaearctic species known from Austria, Croatia, France, Georgia, Germany, Greece, Italy, Portugal, Romania, Slovakia, Slovenia, Spain, Switzerland and Hungary (Schillhammer & Schuh 2004, Rücker 2018). In the Czech Republic, a single record from Central Bohemia reported by Míka (2001). New species for Moravia.

Melanophthalma (***Melanophthalma***) ***parvicollis*** (Mannerheim, 1844). Moravia mer.: Pavlice (7061), 7.ix.2017, 1 ♂, sweeping on horse pasture, R. Stejskal leg., det. et coll., P. Průdek revid.; Plenkovice (7061), 13.viii.2017, 2 ♂♂, near Plenkovický rybník pond, R. Stejskal leg., P. Průdek det., coll. R. Stejskal and Jihomoravské muzeum ve Znojmě; Oleksovice env., Oleksovická mokřina Nature Monument (7063), 19.v.2017, 1 ♂, 22.vi.2017, 1 ♂, both spec. R. Stejskal leg. et coll., P. Průdek det.; same locality, 16.viii.2017, 2 ♂♂, R. Stejskal leg., det. et coll., P. Průdek revid.; Vémyslice env., Na Kocourkách Nature Reserve (7063), 8.vii.2018, 1 ♂, R. Stejskal lgt., det. et coll.; Podyjí National Park, Čížov – Klaperův potok brook (7161), 19.vii.2017, 1 ♂, J. Procházka (Moravské zemské muzeum, Brno, Czech Republic) leg. et coll., P. Průdek det.; Podyjí National Park, Havraníky (7161), 3.xi.2017, 1 ♂, 7.vii.2018, 1 ♂, both specimens in mixed forest north of village, R. Stejskal leg., det. et coll., P. Průdek revid.; Podyjí National Park, Hnanice – Devět mlýnů (7161), 2.x.2017, 2 ♂♂, R. Stejskal leg., det. et coll., P. Průdek revid.; Podyjí National Park, Podmolí – Barák forest (7161), 19.vii.2017, 1 ♂, J. Procházka leg. et coll., P. Průdek det.; Podyjí National Park, Podmolí – Zlámaná skála (7161), 23.viii.2017, 1 ♂, 9.vi.2017, 1 ♂, J. Procházka leg. et coll., P. Průdek det.; Znojmo-Dobšice (7162), flood-plain forest, 13.i.2018, 1 ♂, on fungi, R. Stejskal leg. et coll., P. Průdek det.; Znojmo, "Husovy sady" municipal park (7162), 20.vi.2017, 1 ♀, R. Stejskal leg. et coll., P. Průdek det.; Znojmo-Oblekovice (7162), 7.vii.2018, 1 ♂, sifted from heap of semidry weeds and grass near garden allotments, R. Stejskal leg., det. et coll.; Znojmo,

Riegrova ulice street (7162), 10.ix.2017, 1 ♂, on branches in garden, R. Stejskal leg., det. et coll., P. Průdek revid.; Oleksovice env., Oleksovické vřesoviště Nature Monument (7163), 30.vi.2017, 1 ♂, R. Stejskal lgt. et coll., P. Průdek det.; Jevišovka env., Trávní Dvůr Nature Monument (northern part) (7164), 29.x.2018, 1 ♂, R. Stejskal lgt., det. et coll.

In Europe recorded from Bosnia and *Herzegovina*, Croatia, France, Hungary, Italy, Macedonia, Montenegro, Serbia, Ukraine, and in Asia from Cyprus, Iran, Lebanon and Turkey (Rücker 2018). New species for the Czech Republic.

Melanophthalma (***Melanophthalma***) ***rispini*** Rücker et Johnson, 2007. Moravia mer.: Plenkovice (7061), 13.viii.2017, 1 ♂, sweeping near Plenkovický rybník pond, R. Stejskal leg. et coll., P. Průdek det.; Rudlice env., Rudlické kopce Nature Monument (7062), 8.vi.2013, 1 ♂, sweeping, R. Stejskal leg. et coll., P. Průdek det.; Oleksovice env., Oleksovická mokřina Nature Monument (7063), 22.vi.2017, 1 ♂, at UV light, A. Reiter leg., P. Průdek det., coll. Jihomoravské muzeum ve Znojmě; Vémyslice env., Na Kocourkách Nature Reserve (7063), 8.vii.2018, 1 ♂, sweeping, R. Stejskal leg., det. et coll.


Melanophthalma parvicollis and *M. rispini* belong to the "*Melanophthalma taurica* species group", which at present includes eight species that can only be reliably identified to species by examining the male genitalia (Rücker & Johnson 2007). *Melanophthalma taurica* (Mannerheim, 1844), also previously reported from the Czech Republic (e.g. Jelínek 1993), is distributed in Eastern Europe and eastwards to Central Asia. The known distribution of *M. rispini* includes France, Germany, Hungary, Italy, Poland, Slovenia, Ukraine, North Africa and Turkey (Rücker 2018). Rücker & Johnson (2007) provide a distribution map of *M. rispini* including the Czech Republic, but without giving precise data. First definite records for the Czech Republic.

Corticariinae

Corticaria aphictoides Reitter, 1898 (Figs. 1–2). Moravia mer.: Oleksovice env., Oleksovická mokřina Nature Monument (7063), 22.vi.2017, 1 ♀, at UV light, R. Stejskal lgt. et coll., P. Průdek det.; Suchohrdly – Suchohrdelský rybník pond (7064), 12.iii.2007, 1 ♀, A. Reiter leg., R. Stejskal det., coll. Jihomoravské muzeum ve Znojmě; Podyjí National Park, Havraníky (7161), 30.iv.2018, 1 ♀, sweeping in forest near U Rumcajse wetland; Podyjí National Park, Podmolí (7161), 26.vi.2018, 1 ♀, sifting of compost in forest plant nursery, both specimens R. Stejskal leg. et coll., P. Průdek det.

This species was long time misinterpreted as *Corticaria longicornis* Herbst, 1783 which is, however, the valid name for *C. abietorum* Motschulsky, 1867. A newly recognized species was described by Johnson (2007) as *C. porochini*, but was soon synonymised with *C. aphictoides* (Rücker 2013).

Corticaria aphictoides is widely distributed in the Palaearctic and Nearctic regions, in Europe it is known from Austria, Czech Republic, Finland, Germany, Hungary, Italy, Latvia, Netherlands, Poland, Russia, Spain, Sweden and Ukraine (Rücker 2018). From the Czech Republic, it is listed by Rücker (2013) without precise data. Several records from Bohemia have been published (Hamet 2009, Moravec & Rébl 2014, Januš 2016). From Moravia, it is known from only one specimen of the paratype series of *C. porochini*, collected in Šenov near Ostrava in 1965 (Johnson 2007). Confirmed occurrence in Moravia after at least 50 years.



Figs 1–3. *Corticaria aphictoides* Reitter, 1898: 1 – habitus; 2 – aedeagus in ventral view; 3 – *Corticaria peezi* Johnson, 2007, aedeagus in ventral view. Scale = 0.5 mm. All drawings by Wolfgang H. Rücker.
Obr. 1–3. *Corticaria aphictoides* Reitter, 1898: 1 – habitus; 2 – aedeagus ventrálně; 3 – *Corticaria peezi* Johnson, 2007, aedeagus ventrálně. Měřítko = 0,5 mm. Autorem všech kreseb je Wolfgang H. Rücker.

Corticaria peezi Johnson, 2007 (Fig. 3). Moravia mer., Podyjí National Park, Znojmo-Hradiště (7162), 21.iv.2012, 1 ♂, sifting in dry grassland on southern slopes, R. Stejskal leg. et coll., P. Průdek det.

Newly recognized species, for a long time misidentified as *C. weisei* Reitter, 1875, which is a synonym of *C. serrata* (Paykull, 1798) (Johnson 2007). A very rare species known from central, southern and eastern Europe: Austria, Bosnia and *Herzegovina*, Czech Republic, France, Germany, Greece, Hungary, Italy, Romania, and South European Russia (Rücker 2018). Benedikt (2011) published a record from western Bohemia. New species for Moravia.

ACKNOWLEDGEMENTS. My thanks to Pavel Průdek (Brno, Czech Republic) for identification of the material, valuable comments on the manuscript and help with literature. Special thanks also to Wolfgang H. Rücker (Neuwied, Germany) who kindly provided me with the drawings used in this paper. I also thank my friend Jiří Procházka (Brno, Czech Republic) for providing me with his specimens from the Podyjí National Park, and all reviewers for formal and language corrections.

BENEDIKT S. 2011: Fauna brouků (Coleoptera) lokality Bystřina – Lužní potok (Evropsky významná lokalita soustavy Natura 2000). (Beetle (Coleoptera) fauna in the locality Bystřina – Lužní Potok (Site of Community Importance Natura 2000)). *Západočeské Entomologické Listy* **2**: 13–36 (in Czech, English abstract).

HAMET A. 2009: Faunistický průzkum brouků (Coleoptera) nivy Labe v úseku Vysoká nad Labem – Němčice (Česká republika, východní Čechy) provedený v roce 2008. (Faunistic survey of beetles (Coleoptera) in the Labe river floodplain between Vysoká nad Labem and Němčice (Czech Republic, Eastern Bohemia) in 2008). *Klapalekiana* **45**: 33–63 (in Czech, English summary).

JANUŠ J. 2016: Brouci (Coleoptera) chráněné krajinné oblasti a biosférické rezervace Křivoklátsko. (Beetles (Coleoptera) of Křivoklátsko Protected Landscape Area and the Biosphere Reserve). *Západočeské Entomologické Listy* **Supplementum 1**: 1–449 (in Czech, English abstract).

JELÍNEK J. 1993: Latridiidae. Pp. 106–107. In: JELÍNEK J. (ed.): Check-list of Czechoslovak Insects IV (Coleoptera). (Seznam československých brouků). *Folia Heyrovskyana* **Supplementum 1**: 3–172 (in English and Czech).

JOHNSON C. 2007: Latridiidae. Pp. 74–80. In: LÖBL I. & SMETANA A. (eds): *Catalogue of Palaearctic Coleoptera. Volume 4. Elateroidea, Derodontoidea, Bostrichoidea, Lymexyloidea, Cleroidea, Cucujoidea*. Apollo Books, Stenstrup, 935 pp.

MÍKA P. 2001: Corticariidae (Coleoptera) středního Polabí (Corticariidae (Coleoptera) from the central Labe lowlands). *Studie a Zprávy Okresního Muzea Praha-východ v Brandýse nad Labem a Staré Boleslavi* **14**: 121–144 (in Czech, English abstract).

MORAVEC P. & RÉBL K. 2014: Výsledky faunistického průzkumu brouků (Coleoptera) na území Chráněné krajinné oblasti a Biosférické rezervace Křivoklátsko (Česká republika). Dodatek II. (Results of faunistic survey of beetles (Coleoptera) in the Křivoklátsko Protected Landscape Area and Biosphere Reserve (Czech Republic). Appendix II). *Elateridarium* **8**: 67–103 (in Czech, English abstract).

RÜCKER W. H. 2013: Wiederbeschreibung und ein neues Synonym zu Corticaria aphictoides Reitter, 1898 (Coleoptera, Latridiidae). *Latridiidae – Mitteilungsblatt für Systematik und Taxonomie der Latridiidae* **10**: 23–26.

RÜCKER W. H. 2018: *Latridiidae und Merophysiidae der West-Paläarktis*. Wolfgang H. Rücker, Neuwied, 676 pp.

RÜCKER W. H. & JOHNSON C. 2007: Revision of Melanophthalma taurica (Mannerheim, 1844) species-group and description of three new species (Coleoptera: Latridiidae). *Latridiidae – Mitteilungsblatt für Systematik und Taxonomie der Latridiidae* 5: 11–24.

SCHILLHAMMER H. & SCHUH R. 2004: Bemerkenswerte Käferfunde aus Österreich (XII) (Coleoptera). *Koleopterologische Rundschau* **74**: 443–445.

Robert STEJSKAL, Administration of the Podyjí National Park, Na Vyhlídce 5,
CZ-669 02 Znojmo, Czech Republic; e-mail: rstejskal@centrum.cz