
FAUNISTIC RECORDS FROM THE CZECH REPUBLIC – 535

Diptera: BombyliidaeAlois ČELECHOVSKÝ¹⁾ & David ČELECHOVSKÝ²⁾

¹⁾ Department of Zoology, Faculty of Science, Palacký University, 17. listopadu 50, CZ-771 46 Olomouc, Czech Republic; e-mail: celechov@prfnw.upol.cz

²⁾ Denisova 299, CZ-751 01 Tovačov, Czech Republic; e-mail: dav.celechov@seznam.cz

Bombylisoma minimum (Scopoli, 1772). Moravia mer.: Podyjí National Park, Znojmo env., Havraníky, Skalky u Havraníků Nature Monument (7162), 48°48'23"N 16°00'32"E, 17.vi.2021, 1 ♂, A. Čelechovský & D. Čelechovský leg., A. Čelechovský det. et coll.; Havraníky, Staré vinice (7161), 48°48'34"N 15°59'43"E, 13.vi.2020, 4 ♂♂, 1.vii.2020, 1 ♂, A. Čelechovský & D. Čelechovský leg., A. Čelechovský det. et coll.; Popice, Havranické vřesoviště National Nature Reserve (7162), 48°49'13"N 16°00'45"E, 14.vii.2020, 1 ♂, 17.vi.2021, 1 ♂, 27.vi.2021, 1 ♂, 1 ♀, A. Čelechovský & D. Čelechovský leg., A. Čelechovský det. et coll.

Widespread species in the Palearctic Region, known mainly from the countries of southern, south-western, and south-eastern Europe (Albania, Bosnia and Herzegovina, Bulgaria, Croatia, France, Greece, Italy, Macedonia, Moldova, Portugal, Romania, Russia, Slovenia, Spain, Turkey, and Ukraine), from some countries of the Middle East (Iran and Syria), and reaching Armenia, Azerbaijan, and Georgia in the east. Occurrence in central Europe is reported from Austria, Hungary, Slovakia, Switzerland, and Poland (Čelechovský & Bosák 2009, Dils & Özbek 2006, Evenhuis & Greathead 1999, 2003). The reported occurrence from the Czech Republic (Greathead 2013) is of unknown origin. This is the first confirmed occurrence from the Czech Republic.

Bombylisoma unicolor (Loew, 1855). Moravia mer.: Podyjí National Park, Znojmo env., Havraníky, Staré vinice (7161), 48°48'34"N 15°59'43"E, 4.vii.2019, 1 ♂, 13.vi.2020, 1 ♂, 1.vii.2020, 1 ♂, 5 ♀♀, 17.vi.2021, 7 ♂♂, 3 ♀♀, 27.vi.2021, 1 ♀, A. Čelechovský & D. Čelechovský leg., A. Čelechovský det. et coll.; Popice, Havranické vřesoviště National Nature Reserve (7162), 48°49'13"N 16°00'45"E, 13.vi.2020, 2 ♂♂, A. Čelechovský & D. Čelechovský leg., A. Čelechovský det. et coll.

Palearctic species reported from some countries of southern, south-western, and south-eastern Europe (Albania, Bulgaria, Croatia, Greece, Italy, Moldova, Portugal, Romania, Russia – southern European territory, Spain, Turkey, and Ukraine), also known from northern Africa (Algeria, Egypt, Libya, and Tunisia), from the Middle East (Iran and Israel), and west Asia (Armenia, Azerbaijan, and Georgia). This species is reported relatively rarely in central Europe, known only from Austria and Hungary (Dils & Özbek 2006, Evenhuis & Greathead 1999, 2003). The species has not yet been reported from Slovakia. New species for the Czech Republic.

REFERENCES

- ČELECHOVSKÝ A. & BOSÁK J. 2009: Bombyliidae Latreille, 1802. In: JEDLIČKA L., KÚDELA M. & ST-LOUKALOVÁ V. (eds): *Checklist of Diptera of the Czech Republic and Slovakia*. Electronic version 2, <http://edvis.sk/diptera2009/families/bombyliidae.htm> (accessed 5 April 2022).
- DILS J. & ÖZBEK H. 2006: Contribution to the Knowledge of the Bombyliidae of Turkey (Diptera). *Linzer Biologische Beiträge* **38** (1): 455–504.
- EVENHUIS N. L. & GREATHEAD D. J. 1999: *World Catalog of Bee Flies (Diptera: Bombyliidae)*. Backhuys Publishers, Leiden, 756 pp.
- EVENHUIS N. L. & GREATHEAD D. J. 2003: World Catalog of Bee Flies (Diptera: Bombyliidae): Corrigenda and Addenda. *Zootaxa* **300**: 1–64.
- GREATHEAD D. J. 2013: Fauna Europaea: Bombyliidae. In: PAPE T. & BEUK P. (eds): *Fauna Europaea: Diptera, Brachycera*. Fauna Europaea version 2017.06. <https://fauna-eu.org> (accessed 5 April 2022).