Klapalekiana, 29: 44, 1993
ISSN 1210-6100

FAUNISTIC RECORDS FROM THE CZECH REPUBLIC - 3

Coleoptera:

Elateridae

Ampedus suecicus Palm, 1976. Bohemia mer., Šumava Mts.: Volary (7049), 6.v.1990, 1 ♀, J.Vávra lgt., det. and coll., under the rind of the log of *Picea excelsa*; Stožec (7148), 28.ix.1980, 1 ♀, J.Chalupek lgt., S.Laibner (Choceň) det. and coll. Moravia bor. or., Beskydy Mts., Staré Hamry (6576), 13.iv.1991, 3 ♂♂, 1 ♀, J.Vávra lgt. and coll.; 2 ♂♂, T.Sitek (Ostrava) lgt. and coll., J.Vávra and T.Sitek det.; 11.v.1991, 1 ♀, T.Sitek lgt., det. and coll., under bark and in the surface layer of fallen red-rotten trunk of *Picea excelsa*, in association with *Ampedus tristis* L., *A. sanguineus* L., *A. aethiops* Lac., *Lacon fasciatus* (L.). Distribution: central and northern Europe, Siberia. New species for the Czech Republic.

Anobiidae

Xestobium austriacum Reitter, 1890. Moravia bor. or., Beskydy Mts.: Ostravice (6476), 16.ix.1990, 2 spec. ex larvae; Lysá hora Mt. (6476), 5.i.1991, 1 spec., 20.vi.1992, 1 spec., J.Vávra lgt., det. and coll.; 6.vi.1987, 4 spec., 16.ix.1990, 2 spec., T.Sitek lgt., det. and coll.; Staré Hamry (6576), 13.iv.1991, 3 spec.; Bumbálka (6576), 22.xi.1992, 1 spec., J.Vávra lgt., det. and coll.; 1 spec., T.Sitek lgt., det. and coll.; Smrk Mt. (6576), 13.v.1990, 1 spec., T.Sitek lgt., det. and coll. Both adults and larvae were found in standing dry trunks of *Abies alba*, in remnants of a primary forest of Beskydy Mts. Distribution: Carpathian species. In Central Europe known from Austria and Ukraine, from Slovakia firstly reported by Gottwald (1965) as *Xestobium declive* Duf. New species for the Czech Republic.

Tenebrionidae

Palorus subdepressus (Wollaston, 1864). Moravia bor. or., Ostrava-Bělský les (6275), 28.viii.1992, 1 spec. It flew to light in the evening, J.Vávra lgt., det. and coll. A cosmopolitan species present throughout the Europe. New species for the Czech Republic.

Corticeus bicoloroides Roubal, 1933. Moravia bor. or., Šilheřovice (6075), 9.ii.1993, 6 spec. in the rotten wood, in semicavity of an old trunk of *Tilia* sp., together with *Crepidophorus mutilatus* Rosenh., *Procraerus tibialis* (Lac.), *Ampedus pomorum* (Herb.), *Eucnemis capucina* Ah., *Mycetophagus populi* F., *Mycetochara linearis* (Illig.) and *Rhyncolus* sp.; Moravia mer., Břeclav-Ladná (7167), 16.ix.1991, 1 spec. in old wood, in semicavity of a *Quercus* sp. trunk, together with *Xestobium rufovillosum* (Deg.) and *Collidium filiforme* F.; all J.Vávra lgt., det. and coll. Formerly described as a variety of *Hypophloeus fasciatus* F., raised to species status by Zábranský (1991). Distribution: central Europe (Germany, Austria, Romania and Slovakia). From the territory of the Czech Republic known from Bohemia (Zábranský 1991). New species for Moravia.

GOTTWALD J. 1965: *Zpr. Čs. Společ. Entomol. ČSAV*, 1(2): 1-5. - ZÁBRANSKÝ P. 1991: *Koleopterol. Rundschau*, 61: 175--180.

Jiří VÁVRA, Regional Museum Ostrava,
Masarykovo nám. 1, CZ-702 00 Ostrava 1, Czech Republic